arXiv:1606.06876v1 [math.AG] 22 Jun 2016

TORSION 1-CYCLES AND BIRATIONAL INVARIANT OF DEGREE 4

SHOUHEI MA

Abstract. We describe the torsion part of the Abel-Jacobi kernel in the Griffiths group of 1-cycles in terms of a birational invariant analogous to the degree 4 unramified cohomology. We also give descriptions, in terms of $\mathcal{H}$-cohomology, of the Griffiths group of 1-cycles, and also of the group of torsion cycles algebraically equivalent to zero of any dimension.

1. Introduction

In this paper we are interested in a connection between algebraic cycles and birational invariant, or more specifically, a torsion subgroup of the Griffiths group of 1-cycles and a certain cohomology group which can be thought of as a "homology counterpart" of the degree 4 unramified cohomology.

Let X be a smooth complex projective variety of dimension d. Let $\mathrm{Griff}_1(X) = \mathrm{CH}_1(X)_{hom}/A_1(X)$ be the Griffiths group of 1-cycles, the group of 1-cycles homologous to zero modulo algebraic equivalence. Let $\mathcal{T}_1(X) \subset \mathrm{Griff}_1(X)$ be the image of torsion 1-cycles which is homologous to zero and whose Abel-Jacobi invariant is also zero. This subtle invariant, measuring deviation of torsion cycles with null Deligne cycle class from $A_1(X)$, was first introduced by Voisin [20]. On the other hand, for an abelian group A, let $\mathcal{H}^q(A)$ be the Zariski sheaf on X associated to the presheaf $U \mapsto H^q(U, A)$. The cohomology group $H^{d-k}(X, \mathcal{H}^d(A))$ is a birational invariant of smooth projective varieties ([11]) which is somewhat analogous to the k-th unramified cohomology. The natural homomorphism

$$H^{d-k}(X, \mathcal{H}^k(\mathbb{Z})) \otimes \mathbb{Q}/\mathbb{Z} \to H^{d-k}(X, \mathcal{H}^k(\mathbb{Q}/\mathbb{Z}))$$

is known to be injective ([11]). We will relate $\mathcal{T}_1(X)$ to this quotient group for $k = 4$.

In order to state our result, recall that the coniveau filtration of $H^k(X, \mathbb{Z})$ is defined by ([12], [13], [5])

$$N^r H^k(X, \mathbb{Z}) = \mathrm{Ker}(H^k(X, \mathbb{Z}) \to \varinjlim_{W \subset X} H^k(X - W, \mathbb{Z})),$$

Supported by Grant-in-Aid for Scientific Research (S) 15H05738.

where W ranges over closed algebraic subsets of X of codimension $\geq r$. We will be concerned with the case $k = 2d-3$, where there can be only one nontrivial filter:

$$0 = N^{d-1}H^{2d-3}(X,\mathbb{Z}) \subset N^{d-2}H^{2d-3}(X,\mathbb{Z}) \subset N^{d-3}H^{2d-3}(X,\mathbb{Z}) = H^{2d-3}(X,\mathbb{Z}).$$

We write

$$\Lambda(X) = H^{2d-3}(X,\mathbb{Z})/N^{d-2}H^{2d-3}(X,\mathbb{Z})$$

and let ${}_{tor}\Lambda(X)$ be its torsion part. The group $\Lambda(X)$ gives a proper analogue of Grothendieck's supérieur cohomological invariant ([12] §9) in degree $2d-3$ (so to say, supérieur homological invariant of degree 3). If his generalized Hodge conjecture ([13]) holds true in degree $2d-3$, then ${}_{tor}\Lambda(X)$ can be thought of as measuring defect of its naive integral version.

Theorem 1.1 (Theorem 4.1)**.** *Let X be a smooth complex projective variety of dimension d. We have a short exact sequence*

$$0 \to {}_{tor}\Lambda(X) \to H^{d-4}(X,\mathcal{H}^d(\mathbb{Q}/\mathbb{Z}))/H^{d-4}(X,\mathcal{H}^d(\mathbb{Z}))\otimes\mathbb{Q}/\mathbb{Z} \to \mathcal{T}_1(X) \to 0.$$

When $\mathrm{CH}_0(X)$ is supported in dimension ≤ 2, e.g., when X is rationally connected, $\Lambda(X)$ is finite, $\mathcal{T}_1(X)$ coincides to $\mathrm{Griff}_1(X)$, and we have

$$0 \to \Lambda(X) \to H^{d-4}(X,\mathcal{H}^d(\mathbb{Q}/\mathbb{Z})) \to \mathrm{Griff}_1(X) \to 0.$$

In §5 we also describe the n-torsion part ${}_nA^p(X)$ of the group $A^p(X)$ of codimension p cycles algebraically equivalent to zero modulo rational equivalence, for any p. We will deduce an exact sequence (Theorem 5.1)

$$0 \to H^{p-1}(X,\mathcal{K}_p)/n \to H^{p-1}(X,\mathcal{H}^p(\mathbb{Z}))/n \to {}_nA^p(X) \to 0,$$

where $\mathcal{K}_p$ is the Zariski sheaf on X associated to the Quillen K-theory (we may also replace it by the Milnor K-theory sheaf).

The interaction between algebraic cycles and birational invariant has been originated from the work of Bloch-Ogus [5], and developed around the nineties by many authors such as Parimala, Colliot-Thélène, Barbieri-Viale, Kahn and others (see, e.g., [18], [8], [1], [2], [14] and references in [8]). In particular, the relationship between degree 3 unramified cohomology $H^3_{nr}(X) = H^0(X,\mathcal{H}^3)$ and codimension 2 cycles was extensively studied. After the Bloch-Kato conjecture was proved by Rost-Voevodsky [19], Colliot-Thélène and Voisin [11] revisited this subject and established a relation between $H^3_{nr}(X,\mathbb{Q}/\mathbb{Z})$ and the integral Hodge defect of codimension 2 cycles. Voisin [20] extended this result to the degree 4 unramified cohomology $H^4_{nr}(X,\mathbb{Q}/\mathbb{Z})$, relating it to $\mathcal{T}^3(X)$. On the other hand, in [11] Colliot-Thélène and Voisin also found a relation between the integral Hodge defect of 1-cycles and $H^{d-3}(X,\mathcal{H}^d(\mathbb{Q}/\mathbb{Z}))$. Theorem 1.1 is thus the "homology" counterpart of Voisin's interpretation of H^4_{nr}, and is the extension of the second theorem of Colliot-Thélène–Voisin to degree 4.

Notation. If A is an abelian group and $n > 0$ a natural number, we denote $A/n = A/nA$ and ${}_nA = \{x \in A | nx = 0\}$. We write ${}_{tor}A$ for the subgroup of torsion elements. If $f : A \to B$ is a homomorphism, ${}_nf : {}_nA \to {}_nB$ and ${}_{tor}f : {}_{tor}A \to {}_{tor}B$ denote its restriction to the respective torsion parts. Whether a sheaf is considered in the Zariski topology or the classical topology will be clear from the context.

2. Preliminaries

Let X be a smooth complex projective variety of dimension d. Let A be an abelian group, which will be one of $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{Q}/\mathbb{Z}$ or $\mathbb{Z}/n$ in the sequel. Let $\mathcal{H}^q(A)$ be the Zariski sheaf on Z associated to the presheaf $U \mapsto H^q(U, A)$ defined by the singular cohomology of Zariski open sets. Bloch-Ogus [5] computed the E_2 page of Grothendieck's coniveau spectral sequence ([12] §10) and obtained a spectral sequence

$$E_2^{p,q} = H^p(X, \mathcal{H}^q(A)) \quad \Rightarrow \quad H^{p+q}(X, A)$$

which converges to the coniveau filtration of $H^k(X, A)$ and which coincides to the Leray spectral sequence for the natural continuous map $X_{cl} \to X_{Zar}$ from the classical topology to the Zariski topology. They showed that

$$H^p(X, \mathcal{H}^q(A)) = 0 \quad \text{for} \quad p > q,$$

$$H^p(X, \mathcal{H}^p(\mathbb{Z})) \simeq \mathrm{NS}^p(X),$$

where $\mathrm{NS}^p(X) = \mathrm{CH}^p(X)/A^p(X)$ is the group of codimension p cycles modulo algebraic equivalence. The edge morphism $\mathrm{NS}^p(X) \to H^{2p}(X, \mathbb{Z})$ equals to the cycle map. Note that $\mathcal{H}^q(A) = 0$ for $q > d$ because smooth affine varieties of dimension d have homotopy type of CW complex of real dimension d. Hence we also have $E_2^{p,q} = 0$ for $q > d$. In this way the Bloch-Ogus spectral sequence is confined to the triangle

$$p \geq 0, \quad q \leq d, \quad p \leq q.$$

The group $E_2^{0,k} = H^0(X, \mathcal{H}^k(A))$ is usually called the k-th unramified cohomology group with coefficients in A, and denoted by $H^k_{nr}(X, A)$. As shown in [8], [2], [11], for smooth projective X, it is birationally invariant, and also stably invariant, namely $H^k_{nr}(X, A) \simeq H^k_{nr}(X \times \mathbb{P}^r, A)$. When $A = \mathbb{Z}/n$, $H^k_{nr}(X, \mathbb{Z}/n)$ can be computed in terms of the Galois cohomology of the function field of X (see [8]), which makes $H^k_{nr}(X, \mathbb{Z}/n)$ an effective obstruction in the Noether problem.

In this paper we are mainly interested in the cohomology group which sits in the position of "homology counterpart" of $H^k_{nr}(X, A)$ in the Bloch-Ogus spectral sequence, that is,

$$H^{d-k}(X, \mathcal{H}^d(A)).$$

It is proven by Colliot-Thélène and Voisin [11] that this group is also birationally invariant for smooth projective X. (This can also be seen using the blow-up formula in [2]). Furthermore,

Lemma 2.1. *The group $H^{d-k}(X, \mathcal{H}^d(A))$, $d = \dim X$, is stably invariant.*

Proof. In [2] Barbieri-Viale proved the general formula

$$H^p(X \times \mathbb{P}^r, \mathcal{H}^q(A)) \simeq \bigoplus_{i=0}^{r} H^{p-i}(X, \mathcal{H}^{q-i}(A)).$$

Putting $q = d + r$ and $p = d + r - k$ where $d = \dim X$, we obtain

$$\begin{aligned} H^{d+r-k}(X \times \mathbb{P}^r, \mathcal{H}^{d+r}(A)) &\simeq \bigoplus_{i=0}^{r} H^{d+r-k-i}(X, \mathcal{H}^{d+r-i}(A)) \\ &= H^{d-k}(X, \mathcal{H}^d(A)) \end{aligned}$$

because over X the sheaf $\mathcal{H}^{q'}$ is zero for $q' > d$. □

Although $H^{d-k}(X, \mathcal{H}^d(A))$ shares a few properties with $H^k_{nr}(X, A)$, these two groups may be rather different, and $H^{d-k}(X, \mathcal{H}^d(A))$ has some defects compared to $H^k_{nr}(X, A)$: for example, $H^{d-k}(X, \mathcal{H}^d(\mathbb{Z}))$ may have torsion; cup product is not defined for $H^{d-k}(X, \mathcal{H}^d(A))$; and currently we do not know how to calculate $H^{d-k}(X, \mathcal{H}^d(\mathbb{Z}/n))$ in terms of the function field of X. It is well-known that $H^{d-1}(X, \mathcal{H}^d(\mathbb{Q}/\mathbb{Z}))$ and $H^{d-2}(X, \mathcal{H}^d(\mathbb{Q}/\mathbb{Z}))$ vanish for rationally connected X (cf. Proposition 4.4). According to a conjecture of Voisin, $H^{d-3}(X, \mathcal{H}^d(\mathbb{Q}/\mathbb{Z}))$ would also vanish for such X (see [21] and references therein). By contrast, there are known many examples of rationally connected X for which $H^2_{nr}(X, \mathbb{Q}/\mathbb{Z}) = {}_{tor}H^3(X, \mathbb{Z})$ or $H^3_{nr}(X, \mathbb{Q}/\mathbb{Z}) = Z^4(X)$ is nontrivial (cf. [8], [11]).

Bloch-Srinivas [6] proved that the Zariski sheaf $\mathcal{H}^3(\mathbb{Z})$ is torsion-free using Merkurjev-Suslin Theorem [16]. This was generalized to $\mathcal{H}^q(\mathbb{Z})$ for arbitrary q by Barbieri-Viale [3] and Colliot-Thélène–Voisin [11], as a consequence of the Bloch-Kato conjecture proved by Rost-Voevodsky [19]. As a result, we have a short exact sequence of Zariski sheaves ([11], [1])

$$0 \to \mathcal{H}^q(\mathbb{Z}) \xrightarrow{n} \mathcal{H}^q(\mathbb{Z}) \to \mathcal{H}^q(\mathbb{Z}/n) \to 0 \tag{2.1}$$

for every $n > 0$. Taking cohomology long exact sequence, we obtain

Proposition 2.2 ([11], [1])**.** *For every $p, q \geq 0$ and $n > 0$ we have an exact sequence*

$$0 \to H^p(X, \mathcal{H}^q(\mathbb{Z}))/n \to H^p(X, \mathcal{H}^q(\mathbb{Z}/n)) \to {}_nH^{p+1}(X, \mathcal{H}^q(\mathbb{Z})) \to 0. \tag{2.2}$$

For example, when $(p, q) = (0, k)$, this gives

$$0 \to H^k_{nr}(X, \mathbb{Z})/n \to H^k_{nr}(X, \mathbb{Z}/n) \to {}_nH^1(X, \mathcal{H}^k(\mathbb{Z})) \to 0.$$

On the mirror edge, namely for $(p,q) = (d-k, d)$, we have

(2.3)
$$0 \to H^{d-k}(X, \mathcal{H}^d(\mathbb{Z}))/n \to H^{d-k}(X, \mathcal{H}^d(\mathbb{Z}/n)) \to {}_nH^{d-(k-1)}(X, \mathcal{H}^d(\mathbb{Z})) \to 0.$$

We remark that the map $H^p(X, \mathcal{H}^q(\mathbb{Z}/n)) \to {}_nH^{p+1}(X, \mathcal{H}^q(\mathbb{Z}))$ in (2.2) is induced from the connecting map in the snake lemma applied to the diagram

(2.4)
$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
\cdots & \longrightarrow & \bigoplus_{X^{(p)}} H^{q-p}(\mathbb{C}(x), \mathbb{Z}) & \longrightarrow & \bigoplus_{X^{(p+1)}} H^{q-p-1}(\mathbb{C}(x), \mathbb{Z}) & \longrightarrow & \cdots \\
 & & \downarrow{\scriptstyle n} & & \downarrow{\scriptstyle n} & & \\
\cdots & \longrightarrow & \bigoplus_{X^{(p)}} H^{q-p}(\mathbb{C}(x), \mathbb{Z}) & \longrightarrow & \bigoplus_{X^{(p+1)}} H^{q-p-1}(\mathbb{C}(x), \mathbb{Z}) & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \\
\cdots & \longrightarrow & \bigoplus_{X^{(p)}} H^{q-p}(\mathbb{C}(x), \mathbb{Z}/n) & \longrightarrow & \bigoplus_{X^{(p+1)}} H^{q-p-1}(\mathbb{C}(x), \mathbb{Z}/n) & \longrightarrow & \cdots \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0. & &
\end{array}$$

Here the horizontals are part of the Bloch-Ogus complexes computing the cohomology of $\mathcal{H}^q(\mathbb{Z})$ and $\mathcal{H}^q(\mathbb{Z}/n)$ (= the q-th row in the E_1 page of the coniveau spectral sequence). The columns are exact by the exactness of (2.1) for smooth Zariski open sets of $\overline{\{x\}}$.

Colliot-Thélène and Voisin [11], [20] studied the sequence (2.2) in the following cases and found connection with algebraic cycles.

- $(p,q) = (0,3)$, with the integral Hodge defect $Z^4(X)$ for codimension 2 cycles [11];
- $(p,q) = (0,4)$, with the torsion Abel-Jacobi kernel $\mathcal{T}^3(X)$ in $\mathrm{Griff}^3(X)$ [20];
- $(p,q) = (d-3, d)$, with the integral Hodge defect $Z_2(X)$ for 1-cycles [11].

In §4 we will study the case $(p,q) = (d-4, d)$ and relate it to (still) 1-cycles.

3. Griffiths group of 1-cycles

Let X be a smooth projective complex variety of dimension d. We consider the Bloch-Ogus spectral sequence in degree $\geq 2d-4$. The situation is somewhat similar (though not identical) to that in degree ≤ 4.



FIGURE 1. Bloch-Ogus spectral sequence in degree $\geq 2d-4$

Proposition 3.1. *We have an exact sequence*

$$H^{2d-4}(X,\mathbb{Z}) \xrightarrow{e_4} H^{d-4}(X,\mathcal{H}^d(\mathbb{Z}))$$

$$\xrightarrow{d_2} H^{d-2}(X,\mathcal{H}^{d-1}(\mathbb{Z})) \to H^{2d-3}(X,\mathbb{Z}) \xrightarrow{e_3} H^{d-3}(X,\mathcal{H}^d(\mathbb{Z})) \tag{3.1}$$

$$\xrightarrow{d_2} \mathrm{NS}_1(X) \xrightarrow{cl} H^{2d-2}(X,\mathbb{Z}) \xrightarrow{e_2} H^{d-2}(X,\mathcal{H}^d(\mathbb{Z})) \to 0.$$

Here cl is the cycle map for 1*-cycles modulo algebraic equivalence, and the kernel of e_i is the coniveau filter $N^{d-i+1}H^{2d-i}(X,\mathbb{Z})$ of $H^{2d-i}(X,\mathbb{Z})$.*

The same exact sequence holds for $\mathbb{Z}/n$-coefficients with $\mathrm{NS}_1(X)$ replaced by $\mathrm{NS}_1(X)/n$.

Proof. As illustrated in Figure 1,

(1) the only nonzero differential from degree $2d-3$ to $2d-2$ is $d_2 : E_2^{d-3,d} \to E_2^{d-1,d-1}$, and there is no nonzero differential from degree $2d-2$ to $2d-1$, so the spectral sequence in degree $2d-2$ degenerates at the E_3 page;
(2) similarly, the only nonzero differential from degree $2d-4$ to $2d-3$ is $d_2 : E_2^{d-4,d} \to E_2^{d-2,d-1}$, so the spectral sequence in degree $2d-3$ degenerates at the E_3 page as well.

The observation (1) gives

$$E_2^{d-3,d} \xrightarrow{d_2} E_2^{d-1,d-1} \xrightarrow{cl} E_\infty^{2d-2} \xrightarrow{e_2} E_2^{d-2,d} \to 0,$$

and (2) gives

$$E_\infty^{2d-4} \xrightarrow{e_4} E_2^{d-4,d} \xrightarrow{d_2} E_2^{d-2,d-1} \to E_\infty^{2d-3} \xrightarrow{e_3} E_2^{d-3,d} \xrightarrow{d_2} E_2^{d-1,d-1}.$$

Join of these two exact sequences gives the desired sequence. Since the Bloch-Ogus spectral sequence converges to the coniveau filtration, the first nontrivial filter $\mathrm{Ker}(e_i) \subset E_\infty^{2d-i}$ is equal to $N^{d-i+1}H^{2d-i}(X,\mathbb{Z})$. □

Colliot-Thélène and Voisin [11] studied the third line of (3.1) and found

$$ {}_nZ_2(X) \simeq {}_nH^{d-2}(X, \mathcal{H}^d(\mathbb{Z})). \tag{3.2} $$

Combining this with (2.3) with $k = 3$, they obtained the exact sequence

$$ 0 \to H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))/n \to H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}/n)) \to {}_nZ_2(X) \to 0. \tag{3.3} $$

On the other hand, the Griffiths group $\mathrm{Griff}_1(X)$ is the kernel of the cycle map *cl* and hence equals to the image of $d_2 : H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to \mathrm{NS}_1(X)$ by (3.1). As in §1, let us denote

$$ \Lambda(X) := H^{2d-3}(X, \mathbb{Z})/N^{d-2}H^{2d-3}(X, \mathbb{Z}), \tag{3.4} $$

which is a finitely generated abelian group and is isomorphic to

$$ \begin{aligned} &\mathrm{Im}(\, e_3 : H^{2d-3}(X, \mathbb{Z}) \to H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))\,) \\ &= \mathrm{Ker}(\, d_2 : H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to \mathrm{NS}_1(X)\,) \end{aligned} $$

by (3.1). Looking the second to third line of (3.1), we thus obtain the following description of $\mathrm{Griff}_1(X)$.

Proposition 3.2. *We have an exact sequence*

$$ 0 \to \Lambda(X) \to H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to \mathrm{Griff}_1(X) \to 0. \tag{3.5} $$

In particular, when $H^{2d-3}(X, \mathbb{Z}) = N^{d-2}H^{2d-3}(X, \mathbb{Z})$ *holds, we have*

$$ \mathrm{Griff}_1(X) \simeq H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})). $$

This is analogous to Bloch-Ogus' description ([5]) of $\mathrm{Griff}^2(X)$

$$ 0 \to H^3(X, \mathbb{Z})/NH^3(X, \mathbb{Z}) \to H^3_{nr}(X, \mathbb{Z}) \to \mathrm{Griff}^2(X) \to 0. $$

It is known that the three terms in (3.5) are all birationally invariant for smooth projective X. For $H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))$ this is proved in [11]; for the other two terms, this results from the corresponding blow-up formulae:

$$ \begin{aligned} \mathrm{CH}_1(\tilde{X}) &\simeq \mathrm{CH}_1(X) \oplus \mathrm{CH}_0(Y), \\ H^{2d-3}(\tilde{X}, \mathbb{Z}) &\simeq H^{2d-3}(X, \mathbb{Z}) \oplus H^{2e-1}(Y, \mathbb{Z}), \end{aligned} $$

where $\tilde{X} \to X$ is the blow-up along a smooth subvariety $Y \subset X$ of dimension e.

Corollary 3.3. *The group* $\mathrm{Griff}_1(X)$ *is finitely generated if and only if* $H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))$ *is.*

Bloch-Srinivas [6] proved that when $\mathrm{CH}_0(X)$ is supported in dimension ≤ 2, $\mathrm{Griff}_1(X)$ is a torsion group. Hence in that case, $\mathrm{Griff}_1(X)$ is finite if and only if $H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))$ is. By the way, this result of Bloch-Srinivas can also be obtained from (3.5) and the fact that $H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))$ is torsion in that case ([11]), but this reasoning will be roundabout.

The assumption $H^{2d-3}(X,\mathbb{Z}) = N^{d-2}H^{2d-3}(X,\mathbb{Z})$ in the second statement of Proposition 3.2 is strong. For example, it holds for complete intersections of dimension $d \geq 4$ because in that case $H^{2d-3}(X,\mathbb{Z})$ is zero. When this assumption holds, we have

$$0 \to \mathrm{Griff}_1(X)/n \to H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}/n)) \to {}_nZ_2(X) \to 0$$

by Colliot-Thélène–Voisin's exact sequence (3.3).

Taking the $\mathrm{Ext}(\mathbb{Z}/n, \cdot)$ long exact sequence for (3.5) (or equivalently, applying the snake lemma to the multiplication by n on (3.5)), we obtain a description of the torsion part of $\mathrm{Griff}_1(X)$.

Proposition 3.4. *We have exact sequences*

$$\begin{aligned} 0 \to {}_n\Lambda(X) \to {}_nH^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to {}_n\mathrm{Griff}_1(X) \\ \to \Lambda(X)/n \to H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))/n \to \mathrm{Griff}_1(X)/n \to 0, \end{aligned}$$

$$\begin{aligned} 0 \to {}_{tor}\Lambda(X) \to {}_{tor}H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to {}_{tor}\mathrm{Griff}_1(X) \\ \to \Lambda(X) \otimes \mathbb{Q}/\mathbb{Z} \to H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \otimes \mathbb{Q}/\mathbb{Z} \to \mathrm{Griff}_1(X) \otimes \mathbb{Q}/\mathbb{Z} \to 0. \end{aligned}$$

Corollary 3.5. *The group ${}_n\mathrm{Griff}_1(X)$ (resp. $\mathrm{Griff}_1(X)/n$) is finite if and only if ${}_nH^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))$ (resp. $H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))/n$) is.*

In the corollary we may also replace $H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}))/n$ by $H^{d-3}(X, \mathcal{H}^d(\mathbb{Z}/n))$ thanks to the exact sequence (3.3).

4. Torsion Abel-Jacobi kernel in $\mathrm{Griff}_1(X)$

Let $J^{2d-3}(X)$ be the intermediate Jacobian of X in degree $2d-3$. The algebraic part $J^{2d-3}(X)_{alg}$ of $J^{2d-3}(X)$ is defined as the image of the Abel-Jacobi map $\lambda_{alg} : A_1(X) \to J^{2d-3}(X)$ from $A_1(X) \subset \mathrm{CH}_1(X)_{hom}$. This is an abelian variety, which corresponds to the sub $\mathbb{Q}$-Hodge structure $N^{d-2}H^{2d-3}(X,\mathbb{Q})$ of $H^{2d-3}(X,\mathbb{Q})$ of weight 1. Consider the quotient complex torus

$$J^{2d-3}(X)_{tr} := J^{2d-3}(X)/J^{2d-3}(X)_{alg}.$$

The Abel-Jacobi map $\lambda : \mathrm{CH}_1(X)_{hom} \to J^{2d-3}(X)$ from $\mathrm{CH}_1(X)_{hom}$ induces a homomorphism $\lambda_{tr} : \mathrm{Griff}_1(X) \to J^{2d-3}(X)_{tr}$. We consider its restriction to the torsion part

$${}_{tor}\lambda_{tr} : {}_{tor}\mathrm{Griff}_1(X) \to {}_{tor}J^{2d-3}(X)_{tr}.$$

The situation is summarized in the commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & {}_{tor}A_1(X) & \longrightarrow & {}_{tor}\mathrm{CH}_1(X)_{hom} & \longrightarrow & {}_{tor}\mathrm{Griff}_1(X) & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle {}_{tor}\lambda_{alg}} & & \downarrow{\scriptstyle {}_{tor}\lambda} & & \downarrow{\scriptstyle {}_{tor}\lambda_{tr}} & & \\ 0 & \longrightarrow & {}_{tor}J^{2d-3}(X)_{alg} & \longrightarrow & {}_{tor}J^{2d-3}(X) & \longrightarrow & {}_{tor}J^{2d-3}(X)_{tr} & \longrightarrow & 0. \end{array}$$

The two rows are exact by the divisibility of $A_1(X)$ and $J^{2d-3}(X)_{alg}$ respectively. The homomorphism ${}_{tor}\lambda_{alg}$ remains surjective by the divisibility of $\mathrm{Ker}(\lambda_{alg})$ (see [4] Lemma 1.4).

Following Voisin [20], we define

$$\mathcal{T}_1(X) := \mathrm{Ker}({}_{tor}\lambda_{tr}) \subset {}_{tor}\mathrm{Griff}_1(X).$$

As shown by Voisin, $\mathcal{T}_1(X)$ coincides to the image of $\mathrm{Ker}({}_{tor}\lambda)$ in $\mathrm{Griff}_1(X)$, which is the definition stated in §1. This equality can be seen from the snake lemma applied to the above commutative diagram.

Theorem 4.1. *We have an exact sequence*

$$0 \to {}_{tor}\Lambda(X) \to H^{d-4}(X, \mathcal{H}^d(\mathbb{Q}/\mathbb{Z}))/H^{d-4}(X, \mathcal{H}^d(\mathbb{Z})) \otimes \mathbb{Q}/\mathbb{Z} \to \mathcal{T}_1(X) \to 0.$$

If $\mathrm{CH}_0(X)$ *is supported in dimension* ≤ 3*, we have*

$$0 \to {}_{tor}\Lambda(X) \to H^{d-4}(X, \mathcal{H}^d(\mathbb{Q}/\mathbb{Z})) \to \mathcal{T}_1(X) \to 0.$$

If $\mathrm{CH}_0(X)$ *is supported in dimension* ≤ 2*, then* $\Lambda(X)$ *is finite,* $\mathcal{T}_1(X)$ *coincides to* $\mathrm{Griff}_1(X)$*, and hence*

$$0 \to \Lambda(X) \to H^{d-4}(X, \mathcal{H}^d(\mathbb{Q}/\mathbb{Z})) \to \mathrm{Griff}_1(X) \to 0. \tag{4.1}$$

Proof. If $\mathrm{CH}_0(X)$ is supported in dimension ≤ 3, $H^{d-4}(X, \mathcal{H}^d(\mathbb{Z}))$ is torsion by [11]. Hence it is annihilated when tensored with $\mathbb{Q}/\mathbb{Z}$. The finiteness of $\Lambda(X)$ when $\mathrm{CH}_0(X)$ is supported in dimension ≤ 2 follows by the same argument and (3.5). In that case, $\mathrm{Griff}_1(X)$ is torsion by Bloch-Srinivas [6], and also $J^{2d-3}(X)_{tr} = 0$ because $H^{2d-3}(X, \mathbb{Q}) = N^{d-2}H^{2d-3}(X, \mathbb{Q})$, so $\mathcal{T}_1(X)$ coincides with $\mathrm{Griff}_1(X)$. We prove the first assertion.

We apply (2.2) and the universal coefficient theorem to the 2nd to 4th terms of (3.1) and its $\mathbb{Z}/n$-coefficients version. This gives

(4.2)

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H^{d-4}(X, \mathcal{H}^d(\mathbb{Z}))/n & \longrightarrow & H^{d-4}(X, \mathcal{H}^d(\mathbb{Z}/n)) & \xrightarrow{\alpha} & {}_nH^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow{\scriptstyle d_2} & & \downarrow{\scriptstyle -_nd_2} & & \\
0 & \longrightarrow & H^{d-2}(X, \mathcal{H}^{d-1}(\mathbb{Z}))/n & \longrightarrow & H^{d-2}(X, \mathcal{H}^{d-1}(\mathbb{Z}/n)) & \xrightarrow{\alpha} & {}_n\mathrm{NS}_1(X) & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle \psi/n} & & \downarrow{\scriptstyle \psi_n} & & \downarrow{\scriptstyle -_ncl} & & \\
0 & \longrightarrow & H^{2d-3}(X, \mathbb{Z})/n & \longrightarrow & H^{2d-3}(X, \mathbb{Z}/n) & \xrightarrow{\beta} & {}_nH^{2d-2}(X, \mathbb{Z}) & \longrightarrow & 0,
\end{array}$$

where all rows and the middle column are exact, and the other two columns are complex. The right column is (up to sign) restriction of the 5th to 7th terms of (3.1) to the n-torsion parts. Commutativity at the lower right, saying that the connecting map α of (2.4) with $(p, q) = (d-2, d-1)$ is translated to the Bockstein homomorphism β via the edge morphisms of the spectral sequence (up to sign), is essentially proved in [10] Proposition 1 (replace

μ_m by $\mathbb{Z}$, and the Gersten-Quillen complex by the Bloch-Ogus complex with coefficients in $\mathbb{Z}$). Commutativity at the upper right, namely compatibility of the connecting map α of (2.4) with the d_2 differential, holds generally. It can be checked in the following way (we refer to [9] §1 for the notation): (1) consider the multiplication-by-n and reduction-to-$\mathbb{Z}/n$-coefficients on the whole exact couple $(\underset{\rightarrow}{D}^{p,q}, \underset{\rightarrow}{E}^{p,q})$ with which the coniveau spectral sequence started; (2) interpolate

$$\varinjlim_{Z} H^{p+q+1}(X - Z_{p+3}, X - Z_{p+1}) \tag{4.3}$$

into the relevant diagram

$$\begin{array}{ccccccc}
\underset{\rightarrow}{E}^{p,q} & \xrightarrow{k} & \underset{\rightarrow}{D}^{p+1,q} & \xrightarrow{j} & \underset{\rightarrow}{E}^{p+1,q} & \xrightarrow{k} & \underset{\rightarrow}{D}^{p+2,q} \\
 & & \uparrow{\scriptstyle i} & \nearrow & \uparrow & & \uparrow{\scriptstyle i} \\
 & & & & (4.3) & & \\
 & & & \searrow & \uparrow & & \\
\underset{\rightarrow}{D}^{p+2,q-1} & \xrightarrow{j} & \underset{\rightarrow}{E}^{p+2,q-1} & \xrightarrow{k} & \underset{\rightarrow}{D}^{p+3,q-1} & \xrightarrow{j} & \underset{\rightarrow}{E}^{p+3,q-1}
\end{array}$$

(multiplication-by-n and reduction-to-$\mathbb{Z}/n$-coefficients are omitted); and then (3) run diagram chasing. It turns out that the results of $d_2 \circ \alpha$ and $-\alpha \circ d_2$ both come from a common element of (4.3) with $\mathbb{Z}$-coefficients which is to be multiplied by n.

Now we apply the snake lemma to the lower two rows of (4.2). The resulting connecting map $\mathrm{Ker}({}_n cl) \to \mathrm{Coker}(\psi/n)$ is rewritten as

$$\lambda_n : {}_n\mathrm{Griff}_1(X) \to \Lambda(X)/n.$$

Then the upper side of (4.2) induces the commutative diagram

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H^{d-4}(X, \mathcal{H}^d(\mathbb{Z}))/n & \longrightarrow & H^{d-4}(X, \mathcal{H}^d(\mathbb{Z}/n)) & \longrightarrow & {}_nH^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow{\scriptstyle d_2} & & \downarrow{\scriptstyle \pi_n} & & \\
0 & \longrightarrow & \mathrm{Ker}(\psi/n) & \longrightarrow & \mathrm{Ker}(\psi_n) & \longrightarrow & \mathrm{Ker}(\lambda_n) & \longrightarrow & 0.
\end{array}$$

Since the middle vertical is surjective by the exactness of the middle column of (4.2), the right vertical π_n is also surjective. Since π_n is restriction of the map $d_2 : H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to \mathrm{NS}_1(X)$ of (3.1) to the n-torsion part, its kernel is

$${}_n\mathrm{Ker}(\, d_2 : H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to \mathrm{NS}_1(X)\,) \simeq {}_n\Lambda(X).$$

Thus we obtain

$$0 \to {}_n\Lambda(X) \to {}_nH^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to \mathrm{Ker}(\lambda_n) \to 0.$$

Taking direct limit with respect to n, we have

$$0 \to {}_{tor}\Lambda(X) \to {}_{tor}H^{d-3}(X, \mathcal{H}^d(\mathbb{Z})) \to \mathrm{Ker}(\lambda_\infty) \to 0$$

where

$$\lambda_\infty := \varinjlim_n \lambda_n : {}_{tor}\mathrm{Griff}_1(X) \to \Lambda(X) \otimes \mathbb{Q}/\mathbb{Z}.$$

The construction of λ_∞ as a connecting map coincides to that of Voisin's map $cl_{d-1,tors,tr}$ in [20] p.354 (more precisely, its restriction to ${}_{tor}\mathrm{Griff}_1(X)$). By [20] Proposition 4.4, we see that $\lambda_\infty = {}_{tor}\lambda_{tr}$ and so $\mathrm{Ker}(\lambda_\infty) = \mathcal{T}_1(X)$. Finally, our assertion follows by taking direct limit of (2.3) with $k = 4$. □

Remark 4.2. (1) This proof was inspired by the argument of Voisin [20] in the H^4_{nr} case. On the other hand, if we add the present argument to Voisin's proof, we obtain

$$0 \to {}_{tor}(NH^5(X,\mathbb{Z})/N^2H^5(X,\mathbb{Z})) \to H^4_{nr}(X,\mathbb{Q}/\mathbb{Z})/H^4_{nr}(X,\mathbb{Z})\otimes\mathbb{Q}/\mathbb{Z} \to \mathcal{T}^3(X) \to 0.$$

Since $H^5(X,\mathbb{Z})/NH^5(X,\mathbb{Z})$ has no torsion ([11], [3]), the first term is equal to ${}_{tor}(H^5(X,\mathbb{Z})/N^2H^5(X,\mathbb{Z}))$. This gives a refinement of Voisin's result.

(2) It seems plausible that the map λ_n (and hence λ_∞) coincides with the connecting map in Proposition 3.4, but we have not checked this. In this direction Theorem 4.1 would follow more directly.

(3) The sequence (4.1) also follows from (3.5) and the vanishing $H^{d-i}(X, \mathcal{H}^d(\mathbb{Q})) = 0$ for $i = 3, 4$.

In the proof of Theorem 4.1 we also obtained a description of the kernel of the "mod n" transcendental Abel-Jacobi map $\lambda_n : {}_n\mathrm{Griff}_1(X) \to \Lambda(X)/n$.

Corollary 4.3. *We have*

$$0 \to {}_n\Lambda(X) \to H^{d-4}(X, \mathcal{H}^d(\mathbb{Z}/n))/(H^{d-4}(X, \mathcal{H}^d(\mathbb{Z}))/n) \to \mathrm{Ker}(\lambda_n) \to 0.$$

This provides a geometric interpretation of $H^{d-4}(X, \mathcal{H}^d(\mathbb{Z}/n))$ analogous to (3.3). For the sake of completeness let us also describe $H^{d-k}(X, \mathcal{H}^d(\mathbb{Z}/n))$ for $k \leq 2$, which are less intriguing.

Proposition 4.4. *(1) We have exact sequences*

$$0 \to H^{d-2}(X, \mathcal{H}^d(\mathbb{Z}))/n \to H^{d-2}(X, \mathcal{H}^d(\mathbb{Z}/n)) \to {}_nH_1(X,\mathbb{Z}) \to 0, \tag{4.4}$$

$$0 \to H^{d-2}(X, \mathcal{H}^d(\mathbb{Z})) \otimes \mathbb{Q}/\mathbb{Z} \to H^{d-2}(X, \mathcal{H}^d(\mathbb{Q}/\mathbb{Z})) \to {}_{tor}H_1(X,\mathbb{Z}) \to 0. \tag{4.5}$$

When X is rationally connected, all three terms in (4.5) *vanish.*

(2) We have

$$H^{d-1}(X, \mathcal{H}^d(\mathbb{Z}/n)) \simeq H^{d-1}(X, \mathcal{H}^d(\mathbb{Z}))/n \simeq H_1(X,\mathbb{Z})/n.$$

This vanishes for rationally connected X.

Proof. These are consequences of (2.3) with $k = 1, 2$ and the complete degeneration of the Bloch-Ogus spectral sequence in degree $\geq 2d-1$, which gives

$$H^{d-1}(X, \mathcal{H}^d(\mathbb{Z})) \simeq H^{2d-1}(X, \mathbb{Z}) \simeq H_1(X, \mathbb{Z}),$$

$$H^d(X, \mathcal{H}^d(\mathbb{Z})) \simeq H^{2d}(X, \mathbb{Z}) \simeq \mathbb{Z}.$$

When X is rationally connected, it is simply connected and so $H_1(X, \mathbb{Z}) = 0$. □

If Voisin's conjecture (cf. [21]) that $Z_2(X) = 0$ for rationally connected X holds true, we have $H^{d-2}(X, \mathcal{H}^d(\mathbb{Z})) = 0$ for such X by (3.2). Then all terms in (4.4) would vanish as well.

5. Torsion cycles in $A^k(X)$

The previous sections dealt with $\mathrm{Griff}_1(X)$ and its torsion subgroup. In this section, as another application of (2.2), we give a description of the torsion part of $A^p(X)$ in terms of $\mathcal{H}$-cohomology, for arbitrary p. This is independent of §3 and §4. Let $\mathcal{K}_p$ be the Zariski sheaf on X associated to the Quillen K-theory.

Theorem 5.1. *Let X be a smooth complex projective variety. We have exact sequences*

$$0 \to H^{p-1}(X, \mathcal{K}_p)/n \to H^{p-1}(X, \mathcal{H}^p(\mathbb{Z}))/n \to {}_nA^p(X) \to 0,$$

$$0 \to H^{p-1}(X, \mathcal{K}_p) \otimes \mathbb{Q}/\mathbb{Z} \to H^{p-1}(X, \mathcal{H}^p(\mathbb{Z})) \otimes \mathbb{Q}/\mathbb{Z} \to {}_{tor}A^p(X) \to 0.$$

Remark 5.2. (1) As explained in the proof, we may also replace $\mathcal{K}_p$ by the Milnor K-theory sheaf $\mathcal{K}_p^M$. Indeed, we have $H^{p-1}(X, \mathcal{K}_p) \simeq H^{p-1}(X, \mathcal{K}_p^M)$.

(2) When $p = 2$, the second sequence reduces to the famous formula ([16], [17])

$${}_{tor}A^2(X) \simeq H^1(X, \mathcal{H}^2(\mathbb{Z})) \otimes \mathbb{Q}/\mathbb{Z} \simeq NH^3(X, \mathbb{Z}) \otimes \mathbb{Q}/\mathbb{Z} \simeq {}_{tor}J^3(X)_{alg}.$$

Proof. We will combine three short exact sequences. Firstly, the $(p-1, p)$ case of (2.2) gives

$$0 \to H^{p-1}(X, \mathcal{H}^p(\mathbb{Z}))/n \to H^{p-1}(X, \mathcal{H}^p(\mathbb{Z}/n)) \to {}_n\mathrm{NS}^p(X) \to 0. \tag{5.1}$$

As explained at (2.4), the map $H^{p-1}(X, \mathcal{H}^p(\mathbb{Z}/n)) \to {}_n\mathrm{NS}^p(X)$ is induced from the boundary map in the snake lemma for the middle and left columns

of the diagram

(5.2)

$$\begin{array}{ccccc}
\bigoplus_{X^{(p-2)}} H^2(\mathbb{C}(x),\mathbb{Z}) & \longrightarrow & \bigoplus_{X^{(p-1)}} H^1(\mathbb{C}(x),\mathbb{Z}) & \longrightarrow & \bigoplus_{X^{(p)}} \mathbb{Z} \\
\downarrow n & & \downarrow n & & \downarrow n \\
\bigoplus_{X^{(p-2)}} H^2(\mathbb{C}(x),\mathbb{Z}) & \longrightarrow & \bigoplus_{X^{(p-1)}} H^1(\mathbb{C}(x),\mathbb{Z}) & \longrightarrow & \bigoplus_{X^{(p)}} \mathbb{Z} \\
\downarrow & & \downarrow & & \downarrow \\
\bigoplus_{X^{(p-2)}} H^2(\mathbb{C}(x),\mathbb{Z}/n) & \longrightarrow & \bigoplus_{X^{(p-1)}} H^1(\mathbb{C}(x),\mathbb{Z}/n) & \longrightarrow & \bigoplus_{X^{(p)}} \mathbb{Z}/n \\
\downarrow & & \downarrow & & \downarrow \\
0 & & 0 & & 0.
\end{array}$$

Secondly, Colliot-Thélène, Sansuc and Soulé derived an exact sequence ([10] and [7] §3.2)

$$0 \to H^{p-1}(X,\mathcal{K}_p)/n \to H^{p-1}(X,\mathcal{H}^p(\mathbb{Z}/n)) \to {}_n\mathrm{CH}^p(X) \to 0. \tag{5.3}$$

As explained in [10], [7], the map $H^{p-1}(X,\mathcal{H}^p(\mathbb{Z}/n)) \to {}_n\mathrm{CH}^p(X)$ is induced from the boundary map in the snake lemma for the middle and left columns of the diagram

(5.4)

$$\begin{array}{ccccc}
\bigoplus_{X^{(p-2)}} K_2\mathbb{C}(x) & \longrightarrow & \bigoplus_{X^{(p-1)}} \mathbb{C}(x)^\times & \longrightarrow & \bigoplus_{X^{(p)}} \mathbb{Z} \\
\downarrow n & & \downarrow n & & \downarrow n \\
\bigoplus_{X^{(p-2)}} K_2\mathbb{C}(x) & \longrightarrow & \bigoplus_{X^{(p-1)}} \mathbb{C}(x)^\times & \longrightarrow & \bigoplus_{X^{(p)}} \mathbb{Z} \\
\downarrow & & \downarrow & & \downarrow \\
\bigoplus_{X^{(p-2)}} H^2(\mathbb{C}(x),\mathbb{Z}/n) & \longrightarrow & \bigoplus_{X^{(p-1)}} H^1(\mathbb{C}(x),\mathbb{Z}/n) & \longrightarrow & \bigoplus_{X^{(p)}} \mathbb{Z}/n. \\
\downarrow & & \downarrow & & \downarrow \\
0 & & 0 & & 0.
\end{array}$$

The upper two rows come from the Gersten-Quillen resolution of $\mathcal{K}_p$ (we may replace $\mathcal{K}_p$ by $\mathcal{K}_p^M$ because the Gersten resolution has been established also for the Milnor K-theory [15] and $K_iF = K_i^MF$ for $i \le 2$). The columns are exact: the middle comes from the Kummer theory, and the left from the Merkurjev-Suslin theorem [16].

Thirdly, since $A^p(X)$ is divisible, the sequence

$$0 \to {}_nA^p(X) \to {}_n\mathrm{CH}^p(X) \to {}_n\mathrm{NS}^p(X) \to 0 \tag{5.5}$$

remains exact.

Combining the three exact sequences (5.1), (5.3), (5.5), we obtain

$$\begin{array}{ccccccccc}
 & & & & 0 & & & & \\
 & & & & \downarrow & & & & \\
0 & \longrightarrow & H^{p-1}(X,\mathcal{K}_p)/n & = & H^{p-1}(X,\mathcal{K}_p)/n & \longrightarrow & 0 & & \\
 & & & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & H^{p-1}(X,\mathcal{H}^p(\mathbb{Z}))/n & \longrightarrow & H^{p-1}(X,\mathcal{H}^p(\mathbb{Z}/n)) & \longrightarrow & {}_n\mathrm{NS}^p(X) & \longrightarrow & 0 \\
 & & & & \downarrow & & \| & & \\
0 & \longrightarrow & {}_nA^p(X) & \longrightarrow & {}_n\mathrm{CH}^p(X) & \longrightarrow & {}_n\mathrm{NS}^p(X) & \longrightarrow & 0 \\
 & & & & \downarrow & & & & \\
 & & & & 0 & & & &
\end{array}$$

Commutativity at the lower right can be checked by comparing the boundary maps in the snake lemmas for (5.2) and (5.4). They are connected through the map $\mathbb{C}(x)^\times \to H^1(\mathbb{C}(x),\mathbb{Z})$ which is induced from the boundary maps of the exponential sequences on smooth Zariski open sets of $\overline{\{x\}}$. Now diagram chasing (or the snake lemma) induces vertical morphisms on the left column of this diagram, and shows that it is exact. □

By Proposition 3.4 and Proposition 5.1 with $p = d-1$, the torsion part of $\mathrm{CH}_1(X)_{hom}$ is controlled by the degree $2d-3$ terms in the E_2 page of the Bloch-Ogus spectral sequence with $\mathbb{Z}$-coefficients.

References

[1] Barbieri-Viale, L. *Cicli di codimensione 2 su varietà unirazionali complesse.* in "K-Theory (Strasbourg 1992)", Astérisque **226** (1994), 13–41.

[2] Barbieri-Viale, L. *$\mathcal{H}$-cohomologies versus algebraic cycles.* Math. Nachr. **184** (1997), 5–57.

[3] Barbieri-Viale, L. *On the Deligne-Beilinson cohomology sheaves.* Ann. K-Theory **1**, no.1 (2016), 3–17.

[4] Bloch, S. *Lectures on algebraic cycles.* (Second edition) Cambridge University Press, 2010.

[5] Bloch, S.; Ogus, A. *Gersten's conjecture and the homology of schemes.* Ann. Sci. École Norm. Sup. (4) **7** (1974), 181–201.

[6] Bloch, S.; Srinivas, V. *Remarks on correspondences and algebraic cycles.* Amer. J. Math. **105** (1983), no. 5, 1235–1253.

[7] Colliot-Thélène, J.-L. *Cycles algébriques de torsion et K-théorie algébrique.* in "Arithmetic algebraic geometry (Trento, 1991)", 1–49, Lecture Notes in Math., **1553**, Springer, 1993.

[8] Colliot-Thélène, J.-L. *Birational invariants, purity and the Gersten conjecture.* in "K-theory and algebraic geometry: connections with quadratic forms and division

algebras (Santa Barbara, CA, 1992)", 1–64, Proc. Symp. Pure Math., **58**, Part 1, Amer. Math. Soc., 1995.

[9] Colliot-Thélène, J.-L.; Hoobler, R.-T.; Kahn, B. *The Bloch-Ogus-Gabber theorem.* in "Algebraic K-theory (Toronto, ON, 1996)", 31–94, Fields Inst. Commun., **16**, Amer. Math. Soc., 1997.

[10] Colliot-Thélène, J.-L.; Sansuc, J.-J.; Soulé, C. *Torsion dans le groupe de Chow de codimension deux.* Duke Math. J. **50** (1983), no. 3, 763–801.

[11] Colliot-Thélène, J.-L.; Voisin, C. *Cohomologie non ramifiée et conjecture de Hodge entière.* Duke Math. J. **161** (2012), no. 5, 735–801.

[12] Grothendieck, A. *Le groupe de Brauer. III.* in "Dix exposés sur la cohomologie des schémas", 88–188, North-Holland, 1968.

[13] Grothendieck, A. *Hodge's general conjecture is false for trivial reasons.* Topology **8** (1969) 299–303.

[14] Kahn, B. *Applications of weight-two motivic cohomology.* Doc. Math. **1** (1996), No. 17, 395–416.

[15] Kerz, M. *The Gersten conjecture for Milnor K-theory.* Invent. Math. **175** (2009), no. 1, 1–33.

[16] Merkurjev, A. S.; Suslin, A. A. *K-cohomology of Severi-Brauer varieties and the norm residue homomorphism.* Math. USSR-Izv. **21** (1983), no. 2, 307–340.

[17] Murre, J. P. *Un résultat en théorie des cycles algébriques de codimension deux.* C. R. Acad. Sci. Paris Sér. I **296** (1983), no. 23, 981–984.

[18] Parimala, R. *Witt groups of affine three-folds.* Duke Math. J. **57** (1988), no. 3, 947–954.

[19] Voevodsky, V. *On motivic cohomology with $\mathbb{Z}/l$-coefficients.* Ann. of Math. (2) **174** (2011), no. 1, 401–438.

[20] Voisin, C. *Degree 4 unramified cohomology with finite coefficients and torsion codimension 3 cycles.* in "Geometry and arithmetic", 347–368, EMS Ser. Congr. Rep., Eur. Math. Soc., 2012.

[21] Voisin, C. *Remarks on curve classes on rationally connected varieties.* in "A celebration of algebraic geometry", 591–599, Clay Math. Proc., **18**, Amer. Math. Soc., 2013.

Department of Mathematics, Tokyo Institute of Technology, Tokyo 152-8551, Japan
E-mail address: ma@math.titech.ac.jp